

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/55, 15/82, A01H 5/00	A1	(11) International Publication Number: WO 96/21737 (43) International Publication Date: 18 July 1996 (18.07.96)
(21) International Application Number: PCT/EP96/00111 (22) International Filing Date: 11 January 1996 (11.01.96) (30) Priority Data: 195 02 053.7 13 January 1995 (13.01.95) DE (71) Applicant (for all designated States except US): INSTITUT FÜR GENBIOLOGISCHE FORSCHUNG BERLIN GMBH [DE/DE]; Ihnestrasse 63, D-14195 Berlin (DE). (72) Inventors; and (75) Inventors/Applicants (for US only): SONNEWALD, Uwe [DE/DE]; Martha-Brautsch-Strasse 7a, D-06467 Hoym (DE). KOSSMANN, Jens [DE/DE]; Koblenzer Strasse 1, D-10715 Berlin (DE). BOWIEN, Botho [DE/DE]; Romstrasse 54, D-37079 Göttingen (DE). (74) Agent: VOSSIUS & PARTNER; P.O. Box 86 07 67, D-81634 Munich (DE).		(81) Designated States: AU, CA, HU, JP, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: PROCESS AND DNA MOLECULES FOR INCREASING THE PHOTOSYNTHESIS RATE IN PLANTS (57) Abstract The invention describes recombinant DNA molecules that allow expression of a deregulated or unregulated fructose-1,6-bisphosphatase (FBPase) in plant cells. Such expression leads to an increase in the photosynthesis rate and biomass production in photosynthetically active cells. Furthermore, the invention describes transgenic plants that show an increased photosynthesis rate due to the expression of a deregulated or unregulated FBPase.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

Process and DNA molecules for increasing the photosynthesis rate in plants

The present invention relates to a process and DNA molecules for increasing the photosynthesis rate in plants as well as for an increased yield of plants. The photosynthesis rate and/or the yield is increased by the expression of a deregulated or unregulated fructose-1,6-bisphosphatase in the cytosol of transgenic plants. The invention also relates to the plant cells and plants obtainable by this process as well as to the use of DNA sequences coding for proteins having the enzymatic activity of a fructose-1,6-bisphosphatase for the production of plants exhibiting an increased photosynthesis rate. The invention furthermore relates to recombinant DNA molecules leading to the expression of a fructose-1,6-bisphosphatase in plant cells and plants and resulting in an increased photosynthesis rate.


Due to the continuously growing need for food which is a result from the ever-growing world population it is one of the objects of research in the field of biotechnology to try to increase the yields of useful plants. One possibility to attain this object is to genetically engineer the metabolism of plants. Respective targets are, e.g., the primary processes of photosynthesis that result in CO₂ fixation, the transport processes that participate in the distribution of the photoassimilates within the plant, but also the metabolic pathways that lead to the synthesis of storage substances such as starch, proteins or fats.

For example, the expression of a procaryotic asparagine synthetase in plant cells has been described which results in transgenic plants inter alia in an increase in biomass production (EP 0 511 979).

Another proposal has been to express a procaryotic polyphosphate kinase in the cytosol of transgenic plants. Such expression results in potato plants in an increase in yield in terms of tuber weight of up to 30%.

EP-A-0 442 592 describes the expression of an apoplastic invertase in potato plants which leads to a modified yield of transgenic plants so modified.

Further approaches have concentrated on a modification of the activities of enzymes that participate in the synthesis of sucrose, the most important transport metabolite in most plant species. In plants the CO₂ fixed in the course of photosynthesis is transported from the plastids to the cytosol in the form of triosephosphates (glyceraldehyde-3-phosphate and dihydroxyacetone phosphate). In the cytosol the enzyme aldolase forms a molecule of fructose-1,6-bisphosphate by condensation of glyceraldehyde-3-phosphate and dihydroxyacetone phosphate. This molecule is converted into a molecule of fructose-6-phosphate which in turn is the substrate for the synthesis of sucrose phosphate by the enzyme sucrose phosphate synthase according to the equation



The conversion of fructose-1,6-bisphosphate into fructose-6-phosphate is catalyzed by the enzyme fructose-1,6-bisphosphatase (in the following: FBPase; EC 3.1.3.11) which is regulated by various substances. For example, fructose-2,6-bisphosphate and AMP are potent inhibitors of said enzyme. AMP is an allosteric inhibitor, while fructose-2,6-bisphosphate binds to the active center of the enzyme (Ke et al., Proc. Natl. Acad. Sci. USA 86 (1989), 1475-1479; Liu et al., Biochem. Biophys. Res. Comm. 161 (1989), 689-695. Plant cells contain both a cytoplasmatic as well as a chloroplastic FBPase coded for by the nuclear genome. The reverse reaction (conversion of fructose-6-phosphate into fructose-1,6-bisphosphate) is catalyzed by the enzyme

phospho-fructokinase (PFK) using ATP. Said enzyme is activated by fructose-6-phosphate, P_i and fructose-2,6-bisphosphate and inhibited by glyceraldehyde-3-phosphate and dihydroxyacetone phosphate. Besides said enzymes another enzyme is present in plant cells, namely pyrophosphate:fructose-6-phosphate-1-phosphotransferase (PFP) which catalyzes both reactions according to the equation:



So far various attempts have been made to manipulate this step in the synthesis of sucrose such that the amount of CO_2 fixed is increased resulting in an increased biomass production. For example, it has been attempted to increase the production of fructose-1,6-bisphosphate by overexpressing a plant FBPase in the cytosol (Juan et al., Supplement to Plant Physiol., Vol. 105 (1994), 118). However, this does not lead to a measurable increase of sucrose synthesis. Antisense-inhibition of the PFP, too, failed to result in a detectable increase of sucrose synthesis in plant cells (Hajirezaei et al., Planta 192 (1994), 16-30). It has been furthermore attempted to influence the reaction catalyzed by FBPase by modifying the concentration of the allosteric inhibitor fructose-2,6-bisphosphate (Kruger and Scott, Biochemical Society Transactions, Transgenic Plants and Plant Biochemistry 22 (1994), 904-909). However, it has been found that an increase in the fructose-2,6-bisphosphate concentration has no effect on the photosynthesis rate and only a minor effect on the synthesis of starch or sucrose.

The problem underlying the present invention is to provide further processes generally useful in plants that allow an increase of the photosynthesis rate in plants and thus an increase in biomass production and yield.

The problem is solved by the provision of the embodiments characterized in the claims.

The invention relates to recombinant DNA molecules containing

- (a) a promoter functional in plant cells and
- (b) a DNA sequence linked with the promoter in sense orientation which codes for a polypeptide having the enzymatic activity of a fructose-1,6-bisphosphatase, with the polypeptide having the enzymatic activity of a fructose-1,6-bisphosphatase being a deregulated or unregulated enzyme.

It has surprisingly been found that by expression of such DNA molecules in plant cells a dramatic increase in the photosynthesis rate in plants so modified can be achieved vis-à-vis wild type plants. The term "deregulated" means that said enzymes are not regulated in the same manner as the FBPase enzymes normally expressed in plant cells. Specifically, these enzymes are subject to other regulatory mechanisms, i.e., they are not inhibited to the same extent by the inhibitors or activated by the activators which normally inhibit or activate plant FBPases. For example, they are not inhibited by fructose-2,6-bisphosphate or AMP to the same extent as FBPases that are normally present in plants.

The term "unregulated FBPase enzymes" as used in the present invention relates to FBPase enzymes that are not subject to regulatory mechanisms in plant cells, specifically to those that are not regulated by AMP, ATP or fructose-2,6-bisphosphate.

An increased photosynthesis rate means that plants that have been transformed with a DNA molecule according to the invention which leads to the synthesis of a deregulated or unregulated FBPase in the plants exhibit an increased photosynthesis rate vis-à-vis non-transformed plants, preferably a photosynthesis rate that is increased by at

last 10%, particularly a photosynthesis rate that is increased by at least 20%, most preferably a photosynthesis rate that is increased by 30-40%.

The promoter contained in the DNA molecules according to the invention in principle may be any promoter functional in plant cells. The expression of the DNA sequence coding for an unregulated or deregulated FBPase in principle may take place in any tissue of a transformed plant and at any point in time, preferably it takes place in photosynthetically active tissues. An example for an appropriate promoter is the 35S promoter of the cauliflower mosaic virus (Odell et al., Nature 313 (1985), 810-812) which allows constitutive expression in all tissues of a plant. However, promoters may be used that lead to the expression of subsequent sequences only in a certain tissue of the plant, preferably in photosynthetically active tissue (see, e.g., Stockhaus et al., EMBO J. 8 (1989), 2245-2251) or at a point in time determinable by external influences (see, e.g., WO93/07279). Beside the promoter a DNA molecule according to the invention may also contain DNA sequences that guarantee further increase of the transcription, for example so-called enhancer elements, or DNA sequences that are located in the transcribed region and guarantee a more efficient translation of the synthesized RNA into the corresponding protein. Such 5'-nontranslated regions may be obtained from viral genes or suitable eucaryotic genes or may be synthetically produced. They may be homologous or heterologous with respect to the promoter used.

Furthermore, the DNA molecules according to the invention may contain 3'-nontranslated DNA sequences that guarantee transcription termination and polyadenylation of the transcript formed. Such termination signals are known and have been described. They are freely interchangeable. Examples for such termination sequences are the 3'-nontranslated regions including the polyadenylation signal

of the nopaline synthase gene (NOS gene) from agrobacteria, or the 3'-nontranslated regions of the genes of the small subunit of ribulose-1,5-bisphosphate carboxylase (ssRUBISCO).

The DNA sequence coding for a polypeptide having the enzymatic activity of an FBPase may be derived from any organism expressing such enzyme. These DNA sequences are preferably DNA sequences coding for FBPase enzymes which, vis-à-vis the FBPase enzymes occurring in wild type plants, are subject to a modified, preferably a reduced regulation by inhibitors, particularly a reduced allosteric regulation. The enzymes coded for by the sequences may be known, naturally occurring enzymes exhibiting a modified regulation by various substances, or enzymes that have been produced by mutagenesis of known enzymes from bacteria, algae, fungi, animals or plants. Particularly, they may be fragments of such enzymes that still exhibit the enzymatic activity of an FBPase, which, however, are deregulated or unregulated vis-à-vis FBPases that naturally occur in plant cells.

In a preferred embodiment of the present invention the DNA sequence coding for a polypeptide having the enzymatic activity of an FBPase is derived from a procaryotic organism, preferably a bacterial organism. Bacterial FBPases are advantageous in that they are not regulated by fructose-2,6-bisphosphate vis-à-vis plant derived FBPases. Many bacterial FBPases in contrast to the plant and animal derived FBPases are not regulated in their enzymatic activity by AMP. It is preferred to use DNA sequences coding for such FBPases.

In another preferred embodiment the DNA molecules according to the invention contain a DNA sequence from *Alcaligenes eutrophus* coding for a fructose-1,6-bisphosphatase, preferably a DNA sequence exhibiting the coding region depicted under Seq ID No. 1. The FBPase enzyme from

Alcaligenes eutrophus having the amino acid sequence indicated under Seq ID No. 1 in contrast to plant and animal derived FBPase enzymes is not inhibited by AMP (Abdelal and Schlegel, J. Bacteriol. 120 (1974), 304-310). The DNA sequence depicted under Seq ID No. 1 is a chromosomal DNA sequence. Beside said FBPase *Alcaligenes eutrophus* has a FBPase coded for by a plasmid (J. Koßmann; thesis, 1988, Georg-August-Universität, Göttingen, Germany).

Beside the above-mentioned DNA sequence from *Alcaligenes eutrophus* further bacterial DNA sequences are known that code for polypeptides having the enzymatic activity of an FBPase and that may be used to construct the DNA molecules according to the invention due to their properties.

For example, the *cfxF* gene from *Xanthobacter flavus* H4-14 (Meijer et al., J. Gen. Microbiol. 136 (1990), 2225-2230; Meijer et al., Mol. Gen. Genet. 225 (1991), 320-330) as well as the *fbp* gene from *Rhodobacter sphaeroides* (Gibson et al., Biochemistry 29 (1990), 8085-8093; GenEMBL data base: accession no. J02922) have been cloned. The *fbp* gene from *Rhodobacter sphaeroides* is particularly suitable since the FBPase enzyme coded for by said gene is not inhibited by AMP.

Furthermore, the DNA sequence of the *fbp* gene from *Escherichia coli* coding for FBPase is known (Sedivy et al., J. Bacteriol. 158 (1984), 1048-1053; Hamilton et al. Nucl. Acids Res. 16 (1988), 8707; Raines et al., Nucl. Acids Res. 16 (1988), 7931-7942), as well as a mutated FBPase that is insensitive to AMP (Sedivy et al., Proc. Natl. Acad. Sci. USA 83 (1986), 1656-1659).

Furthermore known is a DNA sequence from *Nitrobacter vulgaris* coding for an FBPase (GenEMBL data base: accession no. L22884) and that may also be used to construct the DNA molecules according to the invention.

In another preferred embodiment the DNA molecules according to the invention contain DNA sequences from fungi coding for

an FBPase. DNA sequences coding for FBPase are known from, e.g., *Saccharomyces cerevisiae* and *Schizosaccharomyces pombe* (Rogers et al., J. Biol. Chem. 263 (1988), 6051-6057; GenEMBL data base: accession nos. J03207 and J03213).

In another preferred embodiment the DNA molecules according to the invention contain DNA sequences from animal organisms coding for an FBPase, preferably DNA sequences from mammals. For example, from mammals a cDNA sequence is known which codes for the FBPase from rat liver (El-Maghrabi et al., Proc. Natl. Acad. Sci. USA 85 (1988), 8430-8434) as well as cDNA sequences coding for an FBPase from pig liver and pig kidney (Marcus et al., Proc. Natl. Acad. Sci. USA 79 (1982), 7161-7165; Williams et al., Proc. Natl. Acad. Sci. USA 89 (1992), 3080-3082; Burton et al., Biochem. Biophys. Res. Commun. 192 (1993), 511-517; GenEMBL data base: accession no. M86347). Furthermore known is a cDNA sequence coding for an FBPase from humans (El-Maghrabi, J. Biol. Chem. 268 (1993), 9466-9472; GenEMBL data base: accession nos. M19922 and L10320).

In a further preferred embodiment the DNA molecules according to the invention contain a plant-derived DNA sequence coding for an FBPase. Such sequences are likewise known. For example, Hur et al. (Plant Mol. Biol. 18 (1992), 799-802) describe a cDNA coding for the cytosolic FBPase from spinach. Said enzyme has been extensively examined on the biochemical level (Zimmermann et al., J. Biol. Chem. 253 (1978), 5952-5956; Ladrör et al., Eur. J. Biochem. 189 (1990), 89-94). Raines et al. (Nucl. Acids Res. 16 (1988), 7931-7942) describe a cDNA sequence coding for the chloroplast FBPase from wheat. A genomic DNA sequence coding for said enzyme is also described (Lloyd et al., Mol. Genet. 225 (1991), 209-216). Furthermore known are cDNA sequences coding for FBPases from *Arabidopsis thaliana* (GenEMBL data base: accession no. X58148), *Beta vulgaris* (sugar beet; GenEMBL data base: accession no. M80597),

Brassica napus (GenEMBL data base: accession no. L15303),
Pisum sativum (GenEMBL data base: accession no. X68826),
Spinacia oleracea (GenEMBL data base: accession no. X61690)
and *Solanum tuberosum* (GenEMBL data base: accession no. X76946).

The above-described DNA sequences coding for FBPase enzymes can be used to isolate further DNA sequences from other organisms, employing, e.g., conventional methods such as screening cDNA libraries or genomic libraries with appropriate probes.

DNA sequences coding for FBPase enzymes, which in comparison to FBPases naturally occurring in plant cells are not deregulated or unregulated, can be modified with the help of techniques known to the person skilled in the art such that the proteins coded for are deregulated or unregulated. Thus, the DNA molecules according to the invention may comprise DNA sequences which are derived from DNA sequences from procaryotic, plant or animal organisms or from fungi coding for an FBPase. This fact is explained in more detail in the following.

Apart from DNA sequences coding for FBPase enzymes also FBPase enzymes have been purified, biochemically characterized and partially sequenced from a large number of organisms, e.g., the cytosolic and the chloroplast FBPase from spinach (Zimmermann et al., J. Biol. Chem. 253 (1978), 5952-5956; Ladrör et al., Eur. J. Biochem. 189 (1990), 89-94; Zimmermann et al., Eur. J. Biochem. 70 (1976), 361-367; Soulié et al., Eur. J. Biochem. 195 (1991), 671-678; Marcus and Harrsch, Arch. Biochem. Biophys. 279 (1990), 151-157; Marcus et al., Biochemistry 26 (1987), 7029-7035), the FBPase from maize (Nishizawa and Buchanan, J. Biol. Chem. 256 (1981), 6119-6126), the chloroplast FBPase from wheat (Leegood and Walker, Planta 156 (1982), 449-456), the FBPase from *Synchococcus lopolensis* (Grubling et al., Plant Physiol. (1986), 716-720), from *Polysphondylium pallidum* (Rosen, Arch. Biochem. Biophys. 114 (1966), 31-37), from

rabbit liver (Pontremoli et al., Arch. Biochem. Biophys. 114 (1966), 24-30), from pig (Marcus et al., Proc. Natl. Acad. Sci. USA 79 (1982), 7161-7165), from *Rhodospseudomonas palustris* (Springgate and Stachow, Arch. Biochem. Biophys. 152 (1972), 1-12; Springgate and Stachow, Arch. Biochem. Biophys. 152 (1972), 13-20), from *E. coli* (Fraenkel et al., Arch. Biochem. Biophys. 114 (1966), 4-12) as well as two isoforms from *Nocardia opaca* 1b (Amachi and Bowien, J. Gen. Microbiol. 113 (1979), 347-356).

Furthermore, for the FBPases from pig the crystal structures of the complexes of the enzymes were determined with fructose-6-phosphate, AMP, fructose-2,6-bisphosphate and magnesium (Seaton et al., J. Biol. Chem. 259 (1984), 8915-8916; Ke et al., Proc. Natl. Acad. Sci. USA 87 (1990), 5243-5247; Ke et al., J. Mol. Biol. 212 (1989), 513-539; Ke et al., Proc. Natl. Acad. Sci. USA 88 (1991), 2989-2993; Ke et al., Biochemistry 30 (1991), 4412-4420). In so doing, the binding sites of fructose-6-phosphate and AMP could be identified as well as the amino acid residues interacting with these substances. Furthermore, it has been described for the FBPases from pig that the removal of the nucleotides coding for amino acid residues 1-25 leads to the synthesis of an FBPase that is not inhibited by AMP but retains its catalytic properties (Chatterjee et al., J. Biol. Chem. 260 (1985), 13553-13559). DNA sequences coding for such FBPases are preferably used in the present invention.

Sequence comparisons on the level of the nucleotide sequences of the FBPase genes as well as on the level of the amino acid sequence of the FBPase enzymes have likewise been made in large numbers (Marcus et al., Biochem. Biophys. Res. Comm. 135 (1986), 374-381). The result was that certain domains of the FBPase have been relatively highly conserved even between remotely related organisms (Gibson et al., Biochemistry 29 (1990), 8085-8093; Marcus et al., Proc. Natl. Acad. Sci. USA 85 (1988), 5379-5383; Rogers et al., J. Biol. Chem. 263 (1988), 6051-6057). It could, for instance, be shown that the amino acid residues that form the

catalytic center in the FBPase from pig are highly conserved in the FBPase from *Xanthobacter flavus* (Meijer et al., J. Gen. Microbiol. 136 (1990), 2225-2230).

The sequence comparisons of the amino acid sequence of the FBPase from *Rhodobacter sphaeroides*, too, with the sequences of other FBPase enzymes known so far indicate conserved regions as well as amino acid residues that participate in the catalysis or the regulation of enzyme activity (Gibson et al., Biochemistry 29 (1990), 8085-8093).

The regulation of the FBPase enzymes has likewise been extensively examined and described in detail (Tejwani, Advances in Enzymology, Vol. 54 (1983), 121-194).

Altogether, the data known so far for DNA sequences coding for FBPase enzymes, for amino acid sequences of FBPase enzymes, for crystal structures as well as for the regulatory and for kinetic and biochemical properties of the FBPases known so far give such a detailed picture that it is possible with this information to specifically introduce mutations into available DNA sequences that result in a modified regulation of the enzyme activity of the synthesized protein. As already mentioned above, it is, e.g., possible to remove the inhibition by AMP in the FBPase from pig by deleting the 25 N-terminal amino acids of the enzyme. The catalytic activity of the enzyme is not influenced by this deletion. Due to the high degree of conservation of the FBPase genes it should therefore be possible to evoke an insensitivity to AMP in other FBPase enzymes, too, by deleting a sufficiently long region at the N-terminus of the enzyme.

It is furthermore known for chromosomally or plasmid encoded FBPases from *Alcaligenes eutrophus* that the plasmidarily encoded enzyme has a characteristic ATP binding site that is missing in the chromosomally encoded enzyme. The plasmid encoded FBPase exhibits in its amino acid sequence the motif GQCMAGKS which is missing in the chromosomally encoded FBPase. This sequence has been identified as or is discussed as an ATP binding site for the phosphoribulokinase and many

other enzymes. The detected consensus sequence (GXXXXGKT/S) is completely contained in the above-mentioned sequence. It is possible that this sequence is responsible for the binding of ATP and thus for the inhibition of the enzyme activity by ATP, such as is observed in various bacterial FBPases.

It is therefore possible to screen bacterial DNA sequences coding for FBPases and being inhibited by ATP for the presence of comparable ATP binding sites and to inactivate or remove this ATP binding site by techniques known in molecular biology and to thus produce an enzyme that cannot be inhibited by ATP.

In a similar manner the sensitivity to the inhibitor fructose-2,6-bisphosphate may be modified. The data obtained by X-ray structure analysis for crystals of the FBPase from pig as well as the analysis of various mutants have meanwhile made it possible to characterize the binding site for fructose-2-bisphosphate in the active center of the FBPase. For FBPase from pig, e.g., amino acid residues have been modified by site-directed mutagenesis, which appear to be important for the function of the enzyme due to the structural data obtained (Giroux et al., J. Biol. Chem. 269 (1994), 31404-31409 and the pertaining references). It could be shown that the amino acid arginine 243 of the FBPase from pig kidney participates in the substrate binding as well as in the inhibition by fructose-2,6-bisphosphate. By replacing this amino acid by an alanine residue a functional FBPase enzyme could be produced whose affinity for fructose-2,6-bisphosphate is reduced by a factor of 1,000 as compared to the wild type enzyme whereas affinity for the substrate fructose-1,6-bisphosphate is only reduced by a factor of 10 (Giroux et al., J. Biol. Chem. 269 (1994), 31404-31409). It could be furthermore shown for FBPase from rat liver that removal of a lysine residue in the active center which residue is also essential for the binding of fructose-1,6-bisphosphate and fructose-2,6-bisphosphate, results in an enzyme that possesses an affinity to the inhibitor fructose-

2,6-bisphosphate that is reduced by the factor of 1,000 (El-Maghrabi et al., J. Biol. Chem. 267 (1992), 6526-6530).

Mutagenization of the relevant amino acid residues should therefore also allow the production of mutants that are modified in their control by fructose-2,6-bisphosphate vis-à-vis wild type proteins. Due to the high degree of conservation of the amino acid sequence of the FBPase enzymes, particularly in the area of the active center, it should furthermore be possible to apply the results obtained for mutants of the enzyme of a certain organism to enzymes that are derived from other organisms.

A further possibility of identifying amino acid residues essential for the catalysis as well as for the inhibition by fructose-2,6-bisphosphate is the computer-aided simulation of the molecular structure. Amino acid residues that are identified as being relevant can subsequently be specifically modified by site-directed mutagenesis and mutants can be examined for their properties.

For a particularly efficient increase of the photosynthesis rate or of the synthesis of fructose-6-phosphate from fructose-1,6-bisphosphate FBPase enzymes are used that are subject only to a reduced regulation by the inhibitors of plant FBPase enzymes (deregulated FBPase enzymes), preferably by enzymes that are no longer subject to any regulation (unregulated FBPase enzymes). Their catalytic activity, however, remains largely untouched. The coding regions of FBPase genes from bacteria, fungi, animals or plants can be mutagenized in *E. coli* or any other suitable host according to methods known in the art and can subsequently be analyzed for an increased FBPase activity and the regulatory mechanisms. The introduction of mutations can be performed in a specific manner (e.g., by site-directed mutagenesis) using specific oligonucleotides, or unspecifically. In the case of unspecific mutagenesis there is the possibility of amplifying the respective DNA sequences by polymerase chain reaction in the presence of

Mn²⁺ ions instead of Mg²⁺ ions, where the error rate is increased, or the propagation of the respective DNA molecules in the *E. coli* strain XL1-Red which results in a high error rate during replication of the plasmid DNA introduced into the bacteria.

The mutagenized DNA sequences coding for the FBPase enzymes are subsequently introduced for analysis of the FBPase activity into a suitable host, preferably into an FBPase-deficient *E. coli* strain. An example of such a strain is *E. coli* strain DF657 (Sedivy et al., J. Bacteriol. 158 (1984), 1048-1053). For an identification of clones expressing a functional FBPase enzyme the transformed cells are plated onto minimal medium containing, e.g., glycerol and succinate (each in a concentration of 0.4%) as carbon source. Cells that do not express functional FBPase cannot grow on such a medium. A first pointer to the activity of the expressed FBPase can be the growth rates of transformed viable clones. In order to preclude mutations in the promoter region resulting in an increased FBPase activity, the mutated coding DNA sequences that allow growth on a minimal medium have to be recloned into non-mutagenated vectors and again be screened for FBPase activity (again by complementation of a FBPase deficient *E. coli* strain). Mutants that effect a complementation of an FBPase deficient *E. coli* strain even in the second screening round are used for the analysis of FBPase activity in the presence of various inhibitors and activators.

The respective cells are broken up, and the FBPase activity is detected *in vitro* using an enzymatic test. In such a test the buffer in which the test is performed for analyzing the properties of the FBPase that was mutagenated is chosen such that the pH value and the salt concentrations are in the optimum range. The buffer must furthermore contain the substrate fructose-1,6-bisphosphate (about 1 mM) and MgCl₂ (about 5 mM). If plant and animal FBPases are expressed, an SH protection group reagent such as DTT or β -mercaptoethanol should be present in the buffer. The measurement of the

enzyme activity is based on that two other enzymes, phosphoglucose isomerase and glucose-6-phosphate dehydrogenase from yeast which further react the product of the FBPase reaction, the fructose-6-phosphate, as well as NADP are added to the mixture. The phosphoglucose isomerase transfers fructose-6-phosphate into glucose-6-phosphate which in turn is reacted from glucose-6-phosphate dehydrogenase to give 6-phosphoglucono- δ -lactone while forming NADPH. The increase in NADPH can be photometrically determined by measuring the absorption at 334 nm. This increase also allows to determine the FBPase activity.

By adding various inhibitors (AMP, ATP, fructose-2,6-bisphosphate) the influence of the inhibitors on the enzyme activity of the mutated FBPases can be determined with the enzyme test described above.

By comparing these values with the values for the activity of the non-mutated enzyme suitable mutants can be chosen. DNA sequences coding for the deregulated or unregulated mutated proteins can subsequently be used to construct the DNA molecules according to the invention.

The generation of mutations in FBPase genes as well as the selection of suitable mutants in an FBPase deficient *E. coli* strain can also be carried out as described in Sedivy et al. (Proc. Natl. Acad. Sci. USA 83 (1986), 1656-1659). This process already allowed to isolate an AMP-insensitive FBPase.

According to the invention the deregulated or unregulated FBPase may be located in any desired compartment of the plant cells. In preferred embodiments the deregulated or unregulated FBPase is located in the cytosol or in the plastides of plant cells. Methods to construct DNA molecules which ensure the localization of a desired protein in various compartments of plant cells, namely in the cytosol or the plastides, are well known to the person skilled in the art.

Another subject matter of the present invention are transgenic plant cells that are transformed with an above-described DNA molecule according to the invention, or that are derived from such a transformed cell and contain a recombinant DNA molecule according to the invention, preferably stably integrated into their genome. The transgenic plant cells are preferably photosynthetically active cells.

The transgenic cells according to the invention can be used to regenerate whole transgenic plants.

Therefore, the present invention also relates to transgenic plants containing the transgenic plant cells according to the invention. Expression of a deregulated or unregulated FBPase in the cells of said plants results in an increase in the photosynthesis rate, thereby leading to an increase in biomass production and/or in yield as compared to non-transformed plants.

The transgenic plants according to the invention are preferably produced by introducing a DNA molecule according to the invention into plant cells and regenerating whole plants from the transformed cells.

The transfer of a DNA molecule according to the invention into plant cells is preferably performed using suitable plasmids, particularly plasmids that allow stable integration of the DNA molecule into the genome of transformed plant cells, e.g., of binary plasmids. Suitable plant transformation vectors comprise, e.g., vectors derived from the Ti plasmid of *Agrobacterium tumefaciens*, as well as those vectors described by Herrera-Estrella et al. (Nature 303 (1983), 209), Bevan (Nucl. Acids Res. 12 (1984), 8711-8721), Klee et al. (Bio/Technology 3 (1985), 637-642) and in EP-A2-120 516.

Transformation with the DNA molecules according to the invention is basically possible with cells of all plant

species. Both monocotyledonous and dicotyledonous plants are of interest. For various monocotyledonous and dicotyledonous plants transformation techniques have already been described. Preferably, cells of ornamental or useful plants are transformed. The useful plants are preferably crop plants, particularly cereals (e.g., rye, oats, barley, wheat, maize, rice), potato, rape, pea, sugar beet, soy bean, tobacco, cotton, tomato, etc.

The invention furthermore relates to propagation material of a plant according to the invention, such as seeds, fruit, cuttings, tubers, root stocks, etc. containing the cells according to the invention.

The subject matter of the present invention is furthermore the use of DNA sequences coding for deregulated or unregulated FBPase enzymes for the expression in plant cells, preferably in the cytosol or the plastides, as well as for the production of plants which exhibit an increased photosynthesis rate and/or increased biomass production as compared to wild type plants.

The invention furthermore relates to a process for increasing the photosynthesis rate in plants which comprises the expression of DNA molecules in plant cells which code for a fructose-1,6-bisphosphate which is deregulated or unregulated in comparison to FBPases normally produced in plant cells.

Fig. 1 shows plasmid p35S-FBPase-Ae

- A = fragment A: CaMV 35S promoter, nt 6909-7437
(Franck et al., Cell 21 (1980), 285-294)
- B = fragment B: DNA from *Alcalig nes utrophus* coding for the chromosomally encoded fructose-1,6-bisphosphatase; 1113 bp fragment having the DNA sequence depicted under Seq ID No. 1

orientation towards the promoter: sense
C = fragment C: nt 11748-11939 of the T-DNA of the Ti
plasmid pTiACH5 (Gielen et al., EMBO J. 3 (1984),
835-846)

The examples serve to illustrate the invention.
In the examples, the following techniques are used:

1. Cloning techniques

For the cloning in *E. coli* the vector pUC18 was used.
For the plant transformation the gene constructs were cloned
into the binary vector pBinAR (Höfgen and Willmitzer, Plant
Sci. 66 (1990), 221-230).

2. Bacterial strains

For the pUC vectors and for the pBinAR constructs the *E. coli* strain DH5 α (Bethesda Research Laboratories, Gaithersburgh, USA) was used.

Transformation of the plasmids in the potato plants was carried out by using *Agrobacterium tumefaciens* strain C58C1 pGV2260 (Deblaere et al., Nucl. Acids Res. 13 (1985), 4777-4788).

3. Transformation of *Agrobacterium tumefaciens*

Transfer of the DNA was carried out by direct transformation according to the method by Höfgen and Willmitzer (Nucleic Acids Res. 16 (1988), 9877). The plasmid DNA of transformed *Agrobacteria* was isolated according to the method by Birnboim and Doly (Nucleic Acids Res. 7 (1979), 1513-1523) and subjected to gel electrophoretic analysis after suitable restriction.

4. Transformation of potatoes

Ten small leaves of a potato sterile culture (*Solanum tuberosum* L.cv. Désirée) were wounded with a scalpel and placed in 10 ml MS medium (Murashige and Skoog, *Physiol. Plant.* 15 (1962), 473) containing 2% sucrose which contained 50 μ l of a selectively grown overnight culture of *Agrobacterium tumefaciens*. After gently shaking the mixture for 3-5 minutes it was further incubated in the dark for 2 days. For callus induction the leaves were placed on MS medium containing 1.6% glucose, 5 mg/l naphthyl acetic acid, 0.2 mg/l benzyl aminopurine, 250 mg/l claforan, 50 mg/l kanamycin, and 0.80% Bacto agar. After incubation at 25°C and 3,000 lux for one week the leaves were placed for shoot induction on MS medium containing 1.6% glucose, 1.4 mg/l zeatin ribose, 20 mg/l naphthyl acetic acid, 20 mg/l gibberellic acid, 250 mg/l claforan, 50 mg/l kanamycin and 0.80% Bacto agar.

5. Radioactive labelling of DNA fragments

The DNA fragments were radioactively labelled using a DNA Random Primer Labelling Kit of Boehringer (Germany) according to the manufacturer's information.

6. Northern Blot Analysis

RNA was isolated according to standard techniques from leaf tissue of plants. 50 μ g of RNA were separated in an agarose gel (1.5% agarose, 1 x MEN buffer, 16.6% formaldehyde). The gel was shortly rinsed with water after gel run. The RNA was transferred with 20 x SSC by capillary blot on a Hybond N nylon membrane (Amersham, UK). The membrane was then baked at 80°C for two hrs in vacuo.

The membrane was prehybridized in NSEB buffer at 68°C for 2 hrs and was then hybridized in NSEB buffer at 68°C overnight in the presence of the radioactively labelled probe.

7. Plant cultivation

Potato plants were cultivated in a greenhouse under the following conditions:

Light period	16 hrs at 25,000 lux and 22°C
Dark period	8 hrs at 15°C
Humidity	60%

Media and solutions used

20 x SSC	175.3 g	NaCl
	88.2 g	sodium citrate
	ad	1000 ml with ddH ₂ O
	pH	7.0 with 10 N NaOH

10 x MEN	200 mM	MOPS
	50 mM	sodium acetate
	10 mM	EDTA
	pH	7.0

NSEB buffer	0.25 M	sodium phosphate buffer pH 7.2
	7%	SDS
	1 mM	EDTA
	1%	BSA (wt./vol.)

Example 1

Construction of plasmid p35S-FBPase-Ae and introduction of the plasmid into the genome of potato plants

A DNA fragment of 1136 bp length having the DNA sequence indicated under Seq ID No. 1 was isolated from a suitable plasmid using the restriction endonucleases *NsiI* and *BalI*.

This DNA fragment includes the whole coding region for the chromosomally encoded FBPase from *Alcaligenes eutrophus*. The cohesive ends were filled in using the T4-DNA polymerase and the fragment was inserted into the vector pBinAR (Höfgen and Willmitzer, Plant Sci. 66 (1990), 221-230) which had been linearized with *Sma*I. The vector pBinAR is a derivative of the binary vector Bin19 (Bevan, Nucleic Acids Res. 12 (1984), 8711-8721; commercially available from Clontech Laboratories, Inc., USA).

pBinAR was constructed as follows:

A fragment of 529 bp length comprising nucleotides 6909-7437 of the 35S promoter of the cauliflower mosaic virus (Franck et al., Cell 21 (1980), 285-294) was isolated as *Eco*RI/*Kpn*I fragment from plasmid pDH51 (Pietrzak et al., Nucl. Acids Res. 14, 5857-5868) and ligated between the *Eco*RI and *Kpn*I restriction sites of the polylinker of pBin19, resulting in plasmid pBin19-A.

A fragment of 192 bp length was isolated from plasmid pAGV40 (Herrera-Estrella et al., Nature 303, 209-213) using the restriction endonucleases *Pvu*II and *Hind*III, which fragment comprises the polyadenylation signal of gene 3 of the T-DNA of Ti plasmid pTiACH5 (Gielen et al., EMBO J. 3, 835-846) (nucleotides 11749-11939). After addition of *Sph*I linkers to the *Pvu*I restriction site the fragment was ligated into pBin19-A which had been cleaved with *Sph*I and *Hind*III, resulting in pBinAR.

The DNA fragment was inserted into the vector such that the coding region was in sense-orientation towards the 35S promoter.

The resulting plasmid was called p35S-FBPase-Ae and is depicted in Fig. 1.

Insertion of the DNA fragment results in an expression cassette that is composed of fragments A, B and C as follows (Fig. 1):

Fragment A (529 bp) contains the 35S promoter of the cauliflower mosaic virus (CaMV). The fragment comprises

nucleotides 6909 to 7437 of CaMV (Franck et al., Cell 21 (1980), 285-294).

Fragment B comprises the protein-encoding region of the chromosomally encoded FBPase from *Alcaligenes eutrophus*. This fragment was isolated as *NsiI/BalI* fragment as described above and fused to the 35S promoter in pBinAR in sense orientation.

Fragment C (192 bp) contains the polyadenylation signal of gene 3 of the T-DNA of Ti plasmid pTiACH5 (Gielen et al., EMBO J. 3 (1984), 835-846).

The size of the plasmid p35S-FBPase-Ae is about 12 kb.

Vector p35S-FBPase-Ae was transferred to potato plant cells via *Agrobacterium tumefaciens*-mediated transformation. Intact plants were regenerated from the transferred cells.

Success of the genetic modification of the plants is verified by subjecting the total RNA to a northern blot analysis with respect to the synthesis of an mRNA coding for the FBPase from *A. eutrophus*. Total RNA is isolated from leaves of transformed plants according to standard techniques, separated on an agarose gel, transferred to a nylon membrane and hybridized to a radioactively labelled probe exhibiting the sequence depicted under Seq ID No. 1 or part of said sequence. Successfully transformed plants exhibit a band in northern blot analysis that indicates the specific transcript of the FBPase gene from *Alcaligenes eutrophus*.

Example 2

Analysis of transformed potato plants

Potato plants that had been transformed with the plasmid p35S-FBPase-Ae were examined for their photosynthesis rate as compared to non-transformed plants.

The photosynthesis rates were measured with leaf disks in a leaf disk oxygen electrode (LD2; Hansat ch; Kinks Lynn,

England). The measurement was performed under a saturated CO₂ atmosphere at 20°C as described by Schaewen et al. (EMBO J. 9 (1990), 3033-3044). Light intensity was 550-600 PAR (photosynthetic active radiation).

The results of such a measurement of the photosynthesis rate of plants that were transformed with the plasmid p35S-FBPase-Ae (UF1-7) in comparison with that of non-transformed plants is shown in the following table.

Plant	photosynthesis rate [mmol O ₂ x (m ² x h) ⁻¹]	%
Wild type	48 ± 6.1	100 ± 12.7
UF1-7 (p35S-FBPase-Ae)	67 ± 6.3	140 ± 13.1

For wild type plants ten measurements were performed while for the transformed potato plants UF1-7 five measurements were made.

24

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT:

- (A) NAME: Institut fuer Genbiologische Forschung Berlin GmbH
- (B) STREET: Ihnestr. 63
- (C) CITY: Berlin
- (E) COUNTRY: Germany
- (F) POSTAL CODE (ZIP): 14195
- (G) TELEPHONE: +49 30 83000760
- (H) TELEFAX: +49 30 83000736

(ii) TITLE OF INVENTION: Process and DNA molecules for increasing the photosynthesis rate in plants

(iii) NUMBER OF SEQUENCES: 2

(iv) COMPUTER READABLE FORM:

- (A) MEDIUM TYPE: Floppy disk
- (B) COMPUTER: IBM PC compatible
- (C) OPERATING SYSTEM: PC-DOS/MS-DOS
- (D) SOFTWARE: PatentIn Release #1.0, Version #1.30 (EPO)

(vi) PRIOR APPLICATION DATA:

- (A) APPLICATION NUMBER: DE 19502053.7
- (B) FILING DATE: 13-JAN-1995

(2) INFORMATION FOR SEQ ID NO: 1:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1136 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: *Alcaligenes eutrophus*

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 30..1121

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

ATGCATAGCC AATCTATAGG AGACCTGTC ATG CCT GAA GTC CAA AGG ATG ACC
Met Pro Glu Val Gln Arg Met Thr

53

25

CTG	ACG	CAG	TTC	CTG	ATC	GAG	GAA	CGC	CGC	CGC	TAT	CCG	GAT	GCC	AGC	101
Leu	Thr	Gln	Phe	Leu	Ile	Glu	Glu	Arg	Arg	Arg	Tyr	Pro	Asp	Ala	Ser	
10						15					20					
GGC	GGC	TTC	AAC	GGC	CTG	ATT	CTC	AAC	GTC	GCC	ATG	GCC	TGC	AAG	GAA	149
Gly	Gly	Phe	Asn	Gly	Leu	Ile	Leu	Asn	Val	Ala	Met	Ala	Cys	Lys	Glu	
25					30				35						40	
ATC	CGC	CGC	CGC	GTT	GCC	TTC	GGC	GGC	CTG	GGG	GGC	TTG	CAC	GGC	AAG	197
Ile	Ala	Arg	Ala	Val	Ala	Phe	Gly	Ala	Leu	Gly	Gly	Leu	His	Gly	Lys	
				45					50					55		
GCC	AGC	AAT	CAA	GCC	GGA	GAA	GCA	GGG	GCC	GTC	AAC	GTG	CAG	GGC	GAA	245
Ala	Ser	Asn	Gln	Ala	Gly	Glu	Ala	Gly	Ala	Val	Asn	Val	Gln	Gly	Glu	
			60					65					70			
ATC	CAG	CAG	AAG	CTG	GAC	GTG	CTG	AGC	AAT	ACC	ACC	TTC	CTG	CGC	GTC	293
Ile	Gln	Gln	Lys	Leu	Asp	Val	Leu	Ser	Asn	Thr	Thr	Phe	Leu	Arg	Val	
	75						80						85			
AAC	GAG	TGG	GGC	GGG	TAC	CTG	GCC	GGC	ATG	GGC	TCG	GAG	GAG	ATG	GAG	341
Asn	Glu	Trp	Gly	Gly	Tyr	Leu	Ala	Gly	Met	Ala	Ser	Glu	Glu	Met	Glu	
90						95					100					
CGC	CCT	TAC	CAG	ATC	CCG	GAT	CAC	TAC	CCG	CGC	GGC	AAG	TAC	CTG	CTG	389
Ala	Pro	Tyr	Gln	Ile	Pro	Asp	His	Tyr	Pro	Arg	Gly	Lys	Tyr	Leu	Leu	
105					110					115					120	
GTG	TTC	GAT	CCG	CTC	GAC	GGC	TCA	TCC	AAC	ATC	GAC	GTC	AAT	GTC	TCG	437
Val	Phe	Asp	Pro	Leu	Asp	Gly	Ser	Ser	Asn	Ile	Asp	Val	Asn	Val	Ser	
				125					130					135		
GTG	GGC	AGC	ATC	TTC	TCG	GTG	CTG	CGC	GCG	CCT	GAG	GGC	GCA	AGC	GCC	485
Val	Gly	Ser	Ile	Phe	Ser	Val	Leu	Arg	Ala	Pro	Glu	Gly	Ala	Ser	Ala	
			140					145					150			
GTC	ACC	GAG	CAG	GAT	TTC	CTG	CAG	CCC	GGC	AGC	GCC	CAG	GTG	GCG	GCC	533
Val	Thr	Glu	Gln	Asp	Phe	Leu	Gln	Pro	Gly	Ser	Ala	Gln	Val	Ala	Ala	
			155				160					165				
GGC	TAC	GCG	CTC	TAC	GGT	CCC	ACC	ACC	ATG	CTG	GTG	CTG	ACC	GTG	GGC	581
ly	Tyr	Ala	Leu	Tyr	Gly	Pro	Thr	Thr	Met	Leu	Val	Leu	Thr	Val	Gly	
170						175					180					
AAT	GGC	GTC	AAC	GGC	TTC	ACG	CTC	GAT	CCC	AAC	CTG	GGC	GAG	TTC	TTC	629
Asn	Gly	Val	Asn	Gly	Phe	Thr	Leu	Asp	Pro	Asn	Leu	Gly	Glu	Phe	Phe	
185					190					195					200	
CTC	ACG	CAC	CCC	AAC	CTG	CAG	GTG	CCG	GCC	GAT	ACC	CAG	GAA	TTT	GCC	677
Leu	Thr	His	Pro	Asn	Leu	Gln	Val	Pro	Ala	Asp	Thr	Gln	Glu	Phe	Ala	
				205					210					215		
ATC	AAT	GCG	TCG	AAC	AGC	CGC	TTC	TGG	GAA	CG	CCG	GTG	CAG	CGC	TAC	725
Ile	Asn	Ala	Ser	Asn	Ser	Arg	Ph	Trp	Glu	Ala	Pr	Val	Gln	Arg	Tyr	
			220					225					230			
ATC	GCC	GAG	TGC	ATG	GCC	GGC	AAG	AGC	GGG	CCG	CGC	GGC	AAG	GAT	TTC	773
Ile	Ala	Glu	Cys	Met	Ala	Gly	Lys	Ser	Gly	Pro	Arg	Gly	Lys	Asp	Phe	
			235				240						245			

26

AAT ATG CGC TGG ATC GCG TCG ATG GTG GCC GAG GCG CAC CGC ATC CTG	821
Asn Met Arg Trp Ile Ala Ser Met Val Ala Glu Ala His Arg Ile Leu	
250 255 260	
ATG CGT GGC GGC GTC TTC ATG TAC CCG CGC GAC TCC AAG GAT CCC GCC	869
Met Arg Gly Gly Val Phe Met Tyr Pro Arg Asp Ser Lys Asp Pro Ala	
265 270 275 280	
AAG CCG GGC CGC CTG CGC CTG CTG TAC GAG GCC AAT CCG ATC GCC TTC	917
Lys Pro Gly Arg Leu Arg Leu Leu Tyr Glu Ala Asn Pro Ile Ala Phe	
285 290 295	
CTG ATG GAG CAG GCT GGC GGG CGC GCC AGC ACG GGC CGG CAG ACG CTG	965
Leu Met Glu Gln Ala Gly Gly Arg Ala Ser Thr Gly Arg Gln Thr Leu	
300 305 310	
ATG TCG GTG GCG CCG GGT GCG CTG CAC CAG CGC ATT GGC GTG ATC TTC	1013
Met Ser Val Ala Pro Gly Ala Leu His Gln Arg Ile Gly Val Ile Phe	
315 320 325	
GGC TCG CGC AAT GAA GTG GAA CCG ATC GAG GGC TAC CAC ACC GAC CAG	1061
Gly Ser Arg Asn Glu Val Glu Arg Ile Glu Gly Tyr His Thr Asp Gln	
330 335 340	
ACC GAT CCC GAC CTT CCG AGT CCC CTG TTC AAC GAG CGC AGC CTG TTC	1109
Thr Asp Pro Asp Leu Pro Ser Pro Leu Phe Asn Glu Arg Ser Leu Phe	
345 350 355 360	
CGC GCG TCT GCC TGAGGTGCCT GGCCA	1136
Arg Ala Ser Ala	

(2) INFORMATION FOR SEQ ID NO: 2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 364 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

Met Pro Glu Val Gln Arg Met Thr Leu Thr Gln Phe Leu Ile Glu Glu	
1 5 10 15	
Arg Arg Arg Tyr Pro Asp Ala Ser Gly Gly Phe Asn Gly Leu Ile Leu	
20 25 30	
Asn Val Ala Met Ala Cys Lys Glu Ile Ala Arg Ala Val Ala Phe Gly	
35 40 45	
Ala Leu Gly Gly Leu His Gly Lys Ala Ser Asn Gln Ala Gly Glu Ala	
50 55 60	
Gly Ala Val Asn Val Gln Gly Glu Ile Gln Gln Lys Leu Asp Val Leu	
65 70 75 80	

27

Ser Asn Thr Thr Phe Leu Arg Val Asn Glu Trp Gly Gly Tyr L u Ala
 85 90 95

Gly Met Ala Ser Glu Glu Met Glu Ala Pro Tyr Gln Ile Pro Asp His
 100 105 110

Tyr Pro Arg Gly Lys Tyr Leu Leu Val Phe Asp Pro Leu Asp Gly Ser
 115 120 125

Ser Asn Ile Asp Val Asn Val Ser Val Gly Ser Ile Phe Ser Val Leu
 130 135 140

Arg Ala Pro Glu Gly Ala Ser Ala Val Thr Glu Gln Asp Phe Leu Gln
 145 150 155 160

Pro Gly Ser Ala Gln Val Ala Ala Gly Tyr Ala Leu Tyr Gly Pro Thr
 165 170 175

Thr Met Leu Val Leu Thr Val Gly Asn Gly Val Asn Gly Phe Thr Leu
 180 185 190

Asp Pro Asn Leu Gly Glu Phe Phe Leu Thr His Pro Asn Leu Gln Val
 195 200 205

Pro Ala Asp Thr Gln Glu Phe Ala Ile Asn Ala Ser Asn Ser Arg Phe
 210 215 220

Trp Glu Ala Pro Val Gln Arg Tyr Ile Ala Glu Cys Met Ala Gly Lys
 225 230 235 240

Ser Gly Pro Arg Gly Lys Asp Phe Asn Met Arg Trp Ile Ala Ser Met
 245 250 255

Val Ala Glu Ala His Arg Ile Leu Met Arg Gly Gly Val Phe Met Tyr
 260 265 270

Pro Arg Asp Ser Lys Asp Pro Ala Lys Pro Gly Arg Leu Arg Leu Leu
 275 280 285

Tyr Glu Ala Asn Pro Ile Ala Phe Leu Met Glu Gln Ala Gly Gly Arg
 290 295 300

Ala Ser Thr Gly Arg Gln Thr Leu Met Ser Val Ala Pro Gly Ala Leu
 305 310 315 320

His Gln Arg Ile Gly Val Ile Phe Gly Ser Arg Asn Glu Val Glu Arg
 325 330 335

Ile Glu Gly Tyr His Thr Asp Gln Thr Asp Pro Asp Leu Pro Ser Pro
 340 345 350


Leu Phe Asn Glu Arg Ser Leu Phe Arg Ala S r Ala
 355 360

C l a i m s

1. A recombinant DNA molecule containing
 - (a) a promoter functional in plant cells, and
 - (b) a DNA sequence coding for a polypeptide having the enzymatic activity of a fructose-1,6-bisphosphatase and being linked to the promoter in sense orientation,wherein the polypeptide having the enzymatic activity of a fructose-1,6-bisphosphatase is a deregulated or unregulated enzyme.
2. The DNA molecule according to claim 1, wherein the DNA sequence coding for a polypeptide having the enzymatic activity of a fructose-1,6-bisphosphatase originates from a procaryotic organism or is derived from such a DNA sequence.
3. The DNA molecule according to claim 2, wherein the procaryotic organism is *Alcaligenes eutrophus*.
4. The DNA molecule according to claim 3, wherein the DNA sequence coding for a polypeptide having the enzymatic activity of a fructose-1,6-bisphosphatase has the coding region depicted under Seq ID No. 1.
5. The DNA molecule according to claim 1, wherein the DNA sequence coding for a polypeptide having the enzymatic activity of a fructose-1,6-bisphosphatase originates from a plant or an animal organism or a fungus, or is derived from such a DNA sequence.
6. A transgenic plant cell containing a recombinant DNA molecule according to any of claims 1 to 5.
7. A transgenic plant containing plant cells according to claim 6.

8. The plant according to claim 7 which is an ornamental plant.
9. The plant according to claim 7 which is a useful plant.
10. Propagation material of a plant according to any of claims 7 to 9 containing plant cells according to claim 6.
11. Use of DNA sequences coding for a deregulated or unregulated fructose-1,6-bisphosphatase for the expression in plant cells.
12. The use according to claim 11, wherein the expression of a deregulated or unregulated fructose-1,6-bisphosphatase in the plants leads to an increase of the photosynthesis rate and/or of the biomass production as compared to wild type plants.
13. Process for increasing the photosynthesis rate in plants which comprises the expression of a DNA molecule according to any one of claims 1 to 5 in cells of a plant.

1/1

**Fig. 1**

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP 96/00111

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N15/55 C12N15/82 A01H5/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N A01H

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	PLANT PHYSIOLOGY SUPPLEMENT, vol. 105, no. 1, May 1994, page 118 XP002001923 SAN JUAN, A., ET AL.: "Overexpression of cytosolic fructose-1,6 biphosphatase in transgenic tobacco plants" cited in the application see abstract 636 ---	1-13
A	THE PLANT JOURNAL , vol. 6, no. 5, November 1994, pages 637-650, XP002001924 KOSSMANN, J., ET AL.: "Reduction of the chloroplastic fructose-1,6-bisphosphatase in transgenic potato plants impairs photosynthesis and plant growth" see the whole document --- -/-	1-13

☒ Further documents are listed in the continuation of box C.

☐ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- *&* document member of the same patent family

Date of the actual completion of the international search

3 May 1996

Date of mailing of the international search report

10.05.1996

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+ 31-70) 340-3016

Authorized officer

Maddox, A

INTERNATIONAL SEARCH REPORT

Inter. Appl. No.

PCT/EP 96/00111

C(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>GENE 85:247-252(1989).., XP002001925 KOSSMANN J, ET AL.: "Sequence analysis of the chromosomal and plasmid genes encoding phosphoribulokinase from Alcaligenes eutrophus" see the whole document ---</p>	1-13
A	<p>ARCH MICROBIOL 154 (1). 1990. 85-91., XP000569675 WINDHÖVEL, U., ET AL.: "ON THE OPERON STRUCTURE OF THE CFX GENE CLUSTERS IN ALCALIGENES-EUTROPHUS." see the whole document -----</p>	1-13